



# Induced domain walls of QCD axion, and gravitational waves

March 12, 2026  
Gravitational Waves and the Early Universe  
@KMI Nagoya


**Junseok Lee**

Particle Theory and Cosmology Group, Tohoku U.

Based on *JCAP* 10 (2024) 038 arXiv: 2407.09478  
collaboration with Kai Murai, Fuminobu Takahashi, and Wen Yin

# Axion

The axion is the pseudo-Nambu-Goldstone boson that arises from the spontaneous breaking of global U(1) symmetry.



$$\Phi = \frac{f_\phi}{\sqrt{2}} e^{i\frac{\phi}{f_\phi}}$$


# QCD Axion

QCD axion dynamically solves the strong CP problem.

[Peccei, Quinn '77](#), [Weinberg '78](#), [Wilczek '78](#)


QCD axion obtains a periodic potential by the non-perturbative effect of QCD such that the combination  $\theta + a/f_a$  vanishes at the potential minimum.

$$\mathcal{L} \supset -\frac{\alpha_s}{8\pi} \left( \theta + \frac{a}{f_a} \right) G_{\mu\nu} \tilde{G}^{\mu\nu}$$




# Network of strings

If the U(1) symmetry is spontaneously broken **after** inflation, a network of cosmic strings is formed.



# Collapsing strings and domain walls


If  $N_{\text{DW}} = 1$ , the network of cosmic strings rapidly decays due to the domain wall tension.



Taken from Hiramatsu, Kawasaki, Saikawa, Sekiguchi `12

# Stable string-wall network

If  $N_{\text{DW}} > 1$ , the network of cosmic strings and domain walls is long-lived.




e.g.,  $N_{\text{DW}} = 2$

$\mathcal{O}(1)$  walls in each Hubble horizon


$\sim$  Hubble length

**Axion domain walls generate domain-wall-like structures via mixing with another axion.**

**“Induced domain wall”**



Couple  $\phi$  to gluons:  $\phi G\tilde{G}$



Couple  $\phi$  to gluons:  $\phi G\tilde{G}$

Introduce QCD axion  $a$ :  $aG\tilde{G}$

$$\frac{a_{\text{left}}}{f_a} = -\theta_{\text{left}}$$

$$\theta = \theta_{\text{left}}$$

$$\phi = \phi_{\text{left}}$$

Domain wall of  $\phi$

$$\phi = \phi_{\text{right}}$$


$$\theta = \theta_{\text{right}}$$

$$\frac{a_{\text{right}}}{f_a} = -\theta_{\text{right}}$$


$x$

Couple  $\phi$  to gluons:  $\phi G\tilde{G}$

Introduce QCD axion  $a$ :  $aG\tilde{G}$



**Domain walls for  $a$  are formed even when  $a$  follows a pre-inflationary scenario.**



# Two axions and mixed initial conditions

We consider two axions

$$\begin{array}{cc}
 \text{Heavy axion} & \text{QCD axion} \\
 \Phi = \frac{f_\phi}{\sqrt{2}} e^{i\frac{\phi}{f_\phi}}, & A = \frac{f_a}{\sqrt{2}} e^{i\frac{a}{f_a}}
 \end{array}$$

with the initial conditions:

- { post-inflationary condition for the heavy axion,
- { pre-inflationary condition for the QCD axion.

# Induced domain walls of QCD axion

JL, Murai, Takahashi, Yin '24

$$\begin{aligned}
 V(a, \phi) &= V_{\text{DW}}(\phi) + V_{\text{mix}}(a, \phi) \\
 &= \Lambda^4 \left[ 1 - \cos \left( 2 \frac{\phi}{f_\phi} \right) \right] + \chi(T) \left[ 1 - \cos \left( \frac{a}{f_a} + \frac{\phi}{f_\phi} \right) \right]
 \end{aligned}$$



$\chi(T) \equiv m_a^2(T) f_a^2$  : topological susceptibility

We assume a hierarchy in the mass and the domain wall tension for simplicity.





$$m_\phi \gg m_{a0} \equiv m_a(T=0)$$

$$m_\phi f_\phi^2 \gg m_{a0} f_a^2$$

$$m_\phi/H = 12.0, m_a/H = 1.2$$


 $\phi/f_\phi$  $a/f_a$ 


$$m_\phi/H = 95.2, m_a/H = 9.5$$


 $\phi/f_\phi$  $a/f_a$ 

# Domain walls must decay


We introduce a potential bias along the heavy axion to destabilize domain walls while preserving the PQ mechanism








# Gravitational wave spectrum



# Gravitational wave spectrum



# Summary

- We found a new type of domain-wall-like structure, called “induced domain walls,” arising from axion mixings.
- Heavy axion domain walls can induce domain walls of the QCD axion even with a pre-inflationary initial condition.
- The QCD axion can be produced from the decay of induced domain walls, and can explain dark matter even for  $f_a \gtrsim 3 \times 10^9 \text{ GeV}$ .
- Gravitational waves from the collapse of heavy-axion domain walls can account for the Pulsar Timing Array results.